

L1551NE - Discovery of a Binary Companion

G. H. Moriarty-Schieven

National Research Council of Canada, Herzberg Institute of Astrophysics, Joint Astronomy Centre, 660 N. A'ohoku Pl., Hilo, HI 96720 (g.moriarty-schieven@jach.hawaii.edu)

J. A. Powers

University of Hawaii, Hilo, HI 96720 (ja_powers@yahoo.com)

H. M. Butner

Submillimeter Telescope Observatory, Steward Observatory, University of Arizona, Tucson, AZ 85721 (hbutner@as.arizona.edu)

P. G. Wannier

Jet Propulsion Laboratory, MS 169-506, 4800 Oak Grove Dr., Pasadena, CA 91109 (peter.wannier@jpl.nasa.gov)

and

Jocelyn Keene

California Institute of Technology, MS 320-47, Pasadena, CA 91125 (jkeene@sirtfweb.jpl.nasa.gov)

ABSTRACT

L1551NE is a very young (class 0 or I) low-mass protostar located close to the well-studied L1551 IRS5. We present here evidence, from 1.3mm continuum interferometric observations at $\sim 1''$ resolution, for a binary companion to L1551NE. The companion, whose 1.3mm flux density is $\sim 1/3$ that of the primary component, is located $1.43''$ (~ 230 A.U. at 160pc) to the southeast. The millimeterwave emission from the primary component may have been just barely resolved, with deconvolved size $\sim 0.82'' \times 0.70''$ ($\sim 131 \times 112$ A.U.). The companion emission was unresolved (< 100 A.U.). The pair is embedded within a flattened circum-binary envelope of size $\sim 5.4'' \times 2.3''$ ($\sim 860 \times 370$ A.U.). The masses of the three components (i.e. from the circumstellar material of the primary star and its companion, and the envelope) are approximately 0.044, 0.014 and 0.023 $M_{\odot}$ respectively.

Subject headings: ISM: individual: L1551NE — stars: formation — stars: binary

1. Introduction

Young T Tauri stars in Taurus have been found to have a high incidence of multiplicity, with the fraction of close ($\lesssim 100$ A.U.) companions found to be 0.40 ± 0.08 (Ghez et al. 1993; Leinert et al. 1993; Simon et al. 1995; Patience et al. 1998), using speckle and occultation techniques. Similar surveys of the Hyades cluster found a smaller fraction of 0.30 ± 0.06 , and a still lower fraction of 0.14 ± 0.03 for G dwarfs in the solar neighborhood (Patience et al. 1998), suggesting an evolutionary effect and/or environmental effects during formation. Deeply embedded class I and class 0 protostars cannot be surveyed using similar techniques, since they are not visible at optical wavelengths and are generally too deeply embedded even at infrared wavelengths. Millimeter/sub-millimeter-wavelength interferometry at sub-arcsecond resolution can resolve close companions if they have circumstellar disks. For example, L1551 IRS5 was recently shown to be a binary based on 7mm interferometric observations at the VLA (Rodríguez et al. 1998). We present here evidence that L1551 NE also has a binary companion.

L1551NE (B1950 $4^h28^m50.5^s +18^\circ02'10''$ (Draper, Warren-Smith & Scarrott 1985)) is a young stellar object in the L1551 molecular cloud, at a distance of 160 pc (Snell 1981). Discovered by Emerson et al. (1984) from IRAS data, it is the second brightest embedded source in the Taurus complex after L1551 IRS5, with $L_{bol} \sim 6L_\odot$. It has a molecular outflow (Moriarty-Schieven, Butner & Wannier 1995). The radial density distribution of L1551NE has been modeled by Barsony & Chandler (1993) from $800\mu m$ images, and by Butner et al. (1995) from $100\mu m$ and $200\mu m$ observations. Both found that the density distribution implied by the radial intensity profile was much shallower than the $n(r) \sim r^{-1.5}$ predicted by the “inside-out” collapse model of Terebey, Shu & Cassen (1984). Moriarty-Schieven, Butner & Wannier (1995) have suggested that L1551 NE may be a class 0 source.

2. Observations

The observations were taken with the Owens Valley Radio Observatory millimeter-wave array 1994 November 12 and 1997 January 17, at a wavelength of 1.3mm (230.65 GHz). The 1994 observations were taken with the array (consisting at the time of six 10m antennae) in the compact “l” configuration (maximum baseline 115m, minimum baseline 30m), and in 1997 the data were taken using the extended “h” configuration (maximum baseline 241.7m, minimum baseline 35m). The effective resolution of the synthesized beam was $\sim 2''$ and $\sim 1''$ for the two configurations.

The observations were made in “snapshot” mode, with a series of ten minute integrations interspersed with observations of other sources and the phase calibrators (0528+134 and 3C84), to yield a total integration time of ~ 1 hour in the “ l ” configuration and ~ 2 hours in the “ h ” configuration. The system temperatures were $\sim 1200\text{K}$ and $\sim 1600\text{K}$ respectively. Neptune was used as the primary flux calibrator. The continuum bandwidth is 1GHz.

The data were reduced using the standard reduction package “mma”, and then exported to AIPS which was used to generate and clean images (using the ‘IMAGR’ task and uniform weighting). The clean box included only a small area covering the main source and the apparent binary companion. The angular resolution of the cleaned image was $1.29'' \times 1.07''$ at position angle -74° .

3. Data

Figure 1 presents a contour image of L1551NE. Clearly seen are the primary source (henceforth source A) at (B1950) $04^h28^m50.552^s$ $18^\circ02'09.85''$, and another weaker source $1.43''$ to the southeast (source B). Emission also appears to extend to the north and to the east of the primary source, and surrounding source B.

To verify that we are seeing real sources and not phase instabilities, we generated images using the same techniques of another source, IRAS 04169+2702, which was observed during the same two days and interspersed with L1551NE. Phase errors would be manifested as “anomalous” sources or structures. No such anomalous sources are seen in the image of IRAS 04169+2702. In addition, we cleaned the “ l ” configuration data separately from the “ h ” configuration data. The high-resolution data clearly had two peaks, while the low-resolution data showed an extended disk-like structure with long axis through the line joining the two sources. Thus we believe that the structure seen here in L1551NE is real.

In Table 1 we present the positions, sizes and flux densities of the sources. The single-dish flux density at 1.35mm is also shown.

4. Discussion

There are three distinct components apparent in the image shown in Figure 1; a brighter, possibly extended source at the field center (source A), a weaker, probably point-like source

(B) 1.43" south-east of A, and diffuse, low-level emission which appears to surround both sources and extend $\sim 2''$ to the northwest and east of A.

Sources A and B were fit with elliptical gaussians using the *AIPS* task JMFIT. The primary source A was found to have a size 1.53"x1.28" (i.e. it may have been slightly resolved with a size $\sim 2\sigma$ larger than the beamsize), peak intensity ~ 0.33 Jy and integrated intensity ~ 0.47 Jy (i.e. $\sim 40\%$ ($\sim 3\sigma$) larger than the peak intensity). If it is slightly resolved (deconvolved size $\sim 0.82'' \times 0.70''$), then its size is $\sim 131 \times 112$ A.U. at a distance of 160pc. Source B is located 1.43" (229 A.U. at 160pc) south-east of A, and its size and intensity are consistent with it being unresolved, i.e. < 100 A.U. We estimate the mass within each source using $M_D = \frac{S_\nu D^2}{B_\nu(T)} \frac{4}{3} \frac{a\rho}{Q_\nu}$, and display these in Table 1. We used the integrated flux densities, assumed a distance of 160 pc (Snell 1981), used dust temperature of 42K (Moriarty-Schieven et al. 1994), and dust emissivities from Hildebrand (1983) $\frac{4}{3} \frac{a\rho}{Q_\nu} = 0.1(\lambda/250)^\beta \text{ g cm}^{-2}$, assuming $\beta = 1$ (Moriarty-Schieven et al. 1994), and have assumed a gas to dust mass ratio of 100. The derived masses are approximately 0.044 and 0.014 $M_\odot$ for the emission from source A and B respectively, and 0.022 $M_\odot$ for the envelope. The total mass for the two sources plus envelope is $\sim 0.078 M_\odot$.

Rodríguez, Anglada & Raga (1995) imaged the L1551NE region at $\lambda 3.5$ cm and found a continuum source located within 1.5σ of our source A. Rodríguez et al attribute the $\lambda 3.5$ cm emission to shocks associated with the outflow (Moriarty-Schieven, Butner & Wannier 1995). Thus source A is the likely origin of the L1551NE outflow. Rodríguez et al. found a possible second source located $\sim 0.6''$ east of A, which is not coincident with our source B, for which no $\lambda 3.5$ cm emission was detected. If their source $0.6''$ east of A is real, this would suggest that L1551NE is at least a triple star system. However, another protostar capable of generating an outflow jet should have had a circumstellar disk large enough to be detected by our observations. Possibly the eastern $\lambda 3.5$ cm source is due to a jet from A but offset from the source, or is a background object.

The single-dish flux density (Butner et al. 2000) found using the 14m JCMT telescope (FWHM $20''$) is not significantly different from the total intensity found in our image of L1551NE (primary beam FWHM $28''$). Thus only a small amount, if any, of the single-dish flux can have been resolved out by the interferometer. However, considerable extended emission was seen at $\sim 850 \mu\text{m}$ by Barsony & Chandler (1993) and Moriarty-Schieven et al. (1999), and at $200 \mu\text{m}$ by Butner et al. (1995). Barsony & Chandler and Butner et al. modeled this extended emission as an envelope whose radial density distribution decreases very slowly with distance from the protostar.

Weak, low-level emission can be seen extending $\sim 1-2''$ to the north and east of source A, and perhaps encompassing source B. This extended structure has a disk-like appearance,

of dimension $\sim 5'' \times 2''$ ($\sim 800 \times 300$ pc with long axis at position angle $\sim -12^\circ$. This is roughly perpendicular to the axis of the conical reflection nebula emanating from L1551NE (Draper et al. 1985; Hodapp 1995), and hence of the molecular outflow (Moriarty-Schieven et al. 1995). This disk-like structure may represent a circum-binary disk.

JAP was supported by a Hawai'i Space Grant College Fellowship which is funded by the NASA Undergraduate Space Grant Fellowship program. The Owens Valley millimeter-wave array is supported by NSF grant AST-96-13717.

REFERENCES

- Barsony, M., & Chandler, C. J. 1993, *ApJ*, 406, L71
- Butner, H. M., Moriarty-Schieven, G. H., Ressler, M. W., & Werner, M. W. 1995, *Ap&SS*, 224, 77
- Butner, H. M., Moriarty-Schieven, G. H., & Wannier, P. G. 2000, in preparation
- Draper, P. W., Warren-Smith, R. F., & Scarrott, S. M. 1985, *MNRAS*, 216, 7P
- Emerson, J. P., Harris, S., Jennings, R. E., Beichman, C. A., Baud, B., Beintema, D. A., Marsden, P. L., & Wesselius, P. R. 1984, *ApJ*, 278, L49
- Ghez, A. M., Neugebauer, G., & Matthews, K. 1993, *AJ*, 106, 2005
- Hildebrand, R. H. 1983, *QJRAS*, 24, 267
- Hodapp, K-W. 1995, *ApJS*, 94, 615
- Leinert, C., Zinnecker, H., Weitzel, N., Christou, J., Ridgway, S. T., Jameson, R., Haas, M., & Lenz, R. 1993, *A&A*, 278, 129
- Moriarty-Schieven, G. H., Butner, H. M., Ohashi, N., & Wannier, P. G. 1999, in preparation
- Moriarty-Schieven, G. H., Butner, H. M., & Wannier, P. G. 1995, *ApJ*, 445, L55
- Moriarty-Schieven, G. H., Wannier, P. G., Keene, J., & Tamura, M. 1994, *ApJ*, 436, 800
- Patience, J., Ghez, A. M., Reid, I. N., Weinberger, A. J., & Matthews, K. 1998, *AJ*, 115, 1972
- Rodríguez, L. F., Anglada, G., & Raga, A. 1995, *ApJ*, 454, 149
- Rodríguez, L. F., D'Alessio, P., Wilner, D. J., Ho, P. T. P., Torrelles, J. M., Curiel, S., Gómez, Y., Lizano, S., Pedlar, A., Cantó, J., & Raga, A. C. 1998, *Nature*, 395, 355
- Simon, M., Ghez, A. M., Leinert, Ch., Cassar, L., Chen, W. P., Howell, R. R., Jameson, R. F., Matthews, K., Neugebauer, G., & Richichi, A. 1995, *ApJ*, 443, 625

Snell, R. L. 1981, ApJS, 45, 121

Terebey, S., Shu, F. H., & Cassen, P. 1984, ApJ, 286, 529

Table 1: Source Parameters

Source A	
Peak Intensity	0.333 ± 0.022 Jy/beam
Integrated Intensity	0.473 ± 0.049 Jy
Position (B1950)	$04^h28^m50.559^s$ $18^\circ02'09.91''$ ($\pm 0.1''$)
Size	$1.53'' \times 1.28'' \pm 0.1''$ P.A. $104^\circ \pm 15^\circ$
“Deconvolved” Size	$0.82'' \times 0.70''$ (131×112 A.U. at 160pc)
Mass ^a	$0.044 M_\odot$
Source B	
Peak Intensity	0.146 ± 0.022 Jy/beam
Integrated Intensity	0.197 ± 0.048 Jy
Position (B1950)	$04^h28^m50.604^s$ $18^\circ02'08.64''$ ($\pm 0.17''$)
Mass ^a	$0.014 M_\odot$
Circumbinary Disk	
Integrated Intensity	0.233 ± 0.048 Jy
Size	$5.51'' (\pm 0.44'') \times 2.59'' (\pm 0.22'')$ P.A. $2^\circ (\pm 14^\circ)$
Deconvolved Size	$5.4'' (\pm 0.3'') \times 2.3'' (\pm 0.2'')$ P.A. $2^\circ (\pm 10^\circ)$
Mass ^a	$0.022 M_\odot$
Total Integrated Intensity	
Integ. Intensity	0.851 ± 0.084 Jy
Mass ^a	$0.079 M_\odot$
Single Dish Intensity^b	
19" FWHM beam	0.83 ± 0.03 Jy
Mass ^a	$0.078 M_\odot$

^aAssuming $T_d=42$ K, $M_g/M_d=100$, $D=160$ pc.

^bFrom Butner et al. (2000). Obtained with 14m JCMT (20" FWHM)).




Fig. 1.— Contour image of L1551NE. Source A is the brighter source at the field center, B is $1.43''$ to the southeast. The two lowest positive contours are at 0.049 and 0.07 Jy/beam, and increment at intervals of 0.035 Jy/beam. The negative contours are -0.049 and -0.07 Jy/beam. RMS noise of the image is 0.023 Jy/beam. The synthesized beam (FWHM) is shown in a box in the lower-right corner of the image.